

BOTOX/DYSPOUR/XEOMIN/JEUVEAU/DAXXIFY NEUROMODULATOR PRE AND POST CARE

PRE-TREATMENT CARE:

- Do NOT consume alcoholic beverages for at least twenty-four (24) hours before treatment to minimize the risk of bruising due to alcohol thinning the blood.
- Discontinue Retin-A for two (2) days before and two (2) days after treatment.

POST-TREATMENT CARE:

- Do NOT manipulate the treated area for 3 hours following treatment. Additionally, avoid facial/laser treatments or microdermabrasion for at least ten days post-injections. Consult your provider if you need more clarification about the timeframe for certain services.
- Some providers suggest engaging in smiling, raising eyebrows, and frowning shortly after treatment to facilitate the distribution of Botox into the targeted muscle.
- The patient should refrain from lying down for 4 hours after receiving Botox to prevent Botox from migrating into the eye area, which could lead to drooping eyelids.
- Results may become noticeable within 4 to 7 days. Additional treatments may be necessary if the desired outcome is not achieved after two weeks. The first touch-up within two weeks is complimentary, with subsequent touch-ups incurring charges based on the amount of product used.
- Do NOT engage in strenuous activities, lifting heavy objects, or vigorous exercise for 6 hours after treatment.
- Do NOT wear hats or any headgear for 4 hours post-treatment.
- If possible, avoid wearing makeup immediately after treatment.